

Improvement of Shelf Life and Quality of Mango (*Mangifera indica*) by Edible Coatings

Kumaresan Marappan^{*1}, Vijai Ananth Arumugam² and Kovvasu Khanchana³

¹ Department of Horticulture, Vels Institute of Science, Technology and Advanced Studies, Pallavaram, Chennai - 600 117, Tamil Nadu, India

² Department of Horticulture, JSA College of Agriculture and Technology, Avatti, Tittakudi, Cuddalore - 606 108, Tamil Nadu, India

³ Horticultural Polytechnic College, Mango Research Station, Dr. Y. S. R. Horticultural University, Nuzvid - 521 201, Andhra Pradesh, India

Received: 29 Feb 2024; Revised accepted: 17 May 2024

Abstract

Mango (*Mangifera indica*), considered to be a member of the Anacardiaceae family, is a significant tropical fruit crop with potential markets. After being harvested, mangoes ripen rapidly, which reduces their potential for handling, storing, and transportation. The experiment on the improvement of shelf life and quality of mango by edible coating was laid out in a randomized design (CRD) with ten treatments and three replications. The treatments consist of T₁-Honey Bee Wax 10%, T₂-Honey Bee Wax 25%, T₃-Honey Bee Wax 30%, T₄-Aloe Vera Gel 10%, T₅-Aloe Vera Gel 25%, T₆-Aloe Vera Gel 30%, T₇-Gelatin 10%, T₈-Gelatin 25%, T₉-Gelatin 30%, and T₁₀-Control. The observations on physiological parameters, biochemical parameters, and organoleptic evaluations were recorded on the 3rd, 6th and 9th days after treatment. The results revealed that the mango treated with T₃ (Honey Bee Wax 25%) significantly reduced the physiological loss in weight (0.19, 0.25, and 0.33%) and maximum TSS of 9.29, 10.88, and 14.14 brix, respectively. The lowest spoilage loss at 6th (0.45) and 9th day (0.69), maximum days taken for shriveling initiation (10.50 days), and maximum shelf life (16.35 days) were recorded when the mango fruits were treated with Honey Bee Wax 25% compared to other treatments. Organoleptic characters like fruit color, pulp color, taste, aroma, and overall acceptability showed the maximum scores in treatment T₃ (Honey bee Wax 25%). Honey bee Wax 25% was found to be better for mango fruits to enhance shelf life and maintain the quality of fruits.

Key words: Mango, Edible coating, Honey bee wax, Shelf life, Quality

Mango (*Mangifera indica*) is a valuable tropical fruit crop with high market potential. Mango fruits are rich in nutrients, minerals, carbohydrates, proteins, and lipids, including vitamins A (beta carotene), B₁, B₂, and C (ascorbic acid), and they have a great flavor and aroma [1]. They are also rich in bioactive components, which make them a fantastic source of antioxidants that can delay the aging process, enhance lung function, and lessen the difficulties associated with diabetes, in addition to lowering the risk of some types of cancer [2]. After being harvested, mango fruits ripen quickly, usually within 3 to 9 days. Long-distance commercial shipment of this fruit is severely restricted by the short time [3]. Fruit may only be stored and transported to a limited extent due to its sensitivity to decay, low temperature tolerance, and general perishability from quick ripening and softening. Mangoes are climacteric fruits that mature quickly after being harvested. This restricts their ability to be handled, stored, and transported. A fruit undergoes an array of color, flavor, and texture changes as it ripens, all of which point to compositional alterations. Achieving the highest possible standard of eating appears to require such a shift. The quality, rate of deterioration, and shelf life of fruits are determined by the aforementioned changes and

their rapidity. After harvesting, production and consumption imbalances and the fruits' easy access to postharvest disease infection cause significant losses. Consequently, interventions that slow down the rate at which ripe mangoes ripen during distribution and promote a supply of high-quality ripe fruits could aid in the expansion of mango markets. Therefore, in order to reduce post-harvest losses and increase shelf life while maintaining quality at a reasonable cost, alternative, appropriate post-harvest methods must be found. The development of edible coatings is currently attracting attention since these technologies ought to be straightforward, widely accessible, ecologically benign, and reasonable with no known negative impacts on human health. At present, consumers are interested in safe and biodegradable edible films or coatings derived from natural bio-based products.

Edible coating has proven to be an effective moisture barrier, halting the evaporation of water. Additionally, it improved the fruits' overall look. Fruit firmness, gloss, texture, and resistance to mechanical damage were all enhanced. This edible covering served as a decent gas barrier and, presumably, prevented some of the vitamin C from being deactivated [4]. The fruit's surface is covered in a thin layer of natural layers

***Correspondence to:** Kumaresan Marappan, E-mail: kummutnau@gmail.com; Tel: +91 9677650324

Citation: Marappan K, Arumugam VA, Khanchana K. 2024. Improvement of shelf life and quality of mango (*Mangifera indica*) by edible coatings. Res. Jr. Agril. Sci. 15(3): 781-784.

created by the edible covering. It does, in fact, act as a barrier against microbial contamination, moisture loss, preservation of textural qualities, preservation of color and volatile flavor chemicals, and slowing down the rate of respiration and transpiration of fruits [5]. The use of beeswax coating showed a beneficial influence on quality maintenance in mango [6] and sweet orange [7], even in ambient settings. The edible covering made of beeswax and coconut oil turned out to be a highly effective method for processing strawberries and apricot fruits as little as possible [8]. Extracts from aloe vera decreased the diffusion and transmission of water. Moreover, fruits' physicochemical properties such as weight loss, hardness, pH, and TSS contents were considerably preserved by coverings with varying concentrations of bee wax. According to Amin *et al.* [9], coating also decreased oxygen permeability, which lessened the oxidation of phenolic components in coated fruits.

Using aloe vera gel as a covering can extend the shelf life of mango fruits after harvest while preserving their quality, which lowers postharvest losses. The findings demonstrated that at both temperatures, concentrations of 50 and 75% aloe considerably extended the evidence of shelf life by reducing the decline in titratable acidity. In these treatments, fruit color and ascorbic acid were likewise preserved for 20 days [10]. The shelf life of edible-coated guavas increased by nine days. It indicates the ability of the biodegradable coating to reduce mass loss during storage time. Furthermore, even after 27 days of storage, the fruit had not yet attained senescence, demonstrating how efficient the coating was in slowing down the ripening process [11]. The guavas were coated with cassava starch and bee wax, which helped to preserve the fruits' quality for an extended period of time by postponing mass and chlorophyll loss and preserving their high-quality levels [12]. According to Neeta *et al.* [13], these findings suggest that using 5% zein and 10% gelatin could be a good substitute for preserving the mango fruit's qualitative qualities and increasing its shelf life while it is being stored.

MATERIALS AND METHODS

The investigation on different edible coatings on shelf life and quality of mango (*Mangifera indica*) was carried out in the pomology laboratory, Department of Horticulture, Adhiparasakthi Horticultural College, Tamil Nadu, during 2022. The details of the materials used and the methods adopted during the course of the investigation are presented below. Mango was obtained from the orchard, Department of Horticulture, Faculty of Horticulture, Adhiparasakthi Horticultural College. Honey Bee Wax was obtained from the Kevin Apiary, Coimbatore district, Tamil Nadu. Aloe vera gel was obtained from the nursery, Department of Horticulture, Adhiparasakthi Horticultural College. Gelatin was obtained from the grocery shop in Kalavai, Ranipet. The treatments consist of T₁-Honey Bee Wax 10%, T₂-Honey Bee Wax 25%, T₃-Honey Bee Wax 30%, T₄-Aloe Vera Gel 10%, T₅-Aloe Vera Gel 25%, T₆-Aloe Vera Gel 30%, T₇-Gelatin 10%, T₈-Gelatin 25%, T₉-Gelatin 30%, and T₁₀-Control. Treatments were applied as post-harvest dips. The fruits were dipped for 5 minutes in honey bee wax (10%, 25%, 30%), aloe vera gel (10%, 25%, 30%), and gelatin (10%, 25%, 30%). The fruits were air-dried for 30 minutes after each treatment. Detailed observations were periodically recorded on the 3rd, 6th, and 9th days of storage for all the parameters. The data were statistically analyzed using the analysis of variance technique for a complete randomized design, as described by Panse and Sukhatme [14]. Each case's proper standard errors of means (S.Em (+)) were determined, and the critical difference (C.D.)

at the percent level of probability was computed to compare the two treatment means in cases where the treatment effects were statistically significant.

Organoleptic test

A group of judges evaluated the fruits' flavors and tastes. As listed under mango were the marks assigned to the various grades.

Details	Scale
Like extremely	9-10
like very much	8-9
Like moderately	7-8
Like slightly	6-7
Neither like nor dislike	5-6
Dislike slightly	4-5
Dislike moderately	3-4
Dislike very much	2-3
Dislike extremely	1-2

RESULTS AND DISCUSSION

Physiological loss in weight

The data pertaining to physiological loss in weight at the 3rd, 6th and 9th days after storage are presented in (Table 1). The fruits treated with honey bee wax 25% significantly recorded the lowest physiological loss in weight of 0.19, 0.25, and 0.33% at the 3rd, 6th, and 9th days after storage, respectively, and the control recorded the maximum physiological loss in weight of 5.13, 10.72, and 16.42%, respectively, at the 3rd, 6th, and 9th days after storage. The reduced physiological loss in honey bee wax may be a result of respiration being inhibited by honey bee wax, altering the membrane's permeability [15]. According to Shahid and Abbasim [16], the beeswax coats extended the shelf life of sweet oranges by lowering the respiration rate of the fruits. Fruit dehydration causes a physiological loss of weight due to changes in the fruit's resistance to water vapor transfer on its surface, respiration rate, and the formation of microscopic fissures connecting its internal and external atmospheres. Honey bee wax could reduce endogenous substrate catabolism during respiration and keep it out of reactions associated with internal breakdown in apples by inhibiting the diffusion of substrate from the vacuole to the cytoplasm and promoting the uptake of sorbitol. Beeswax has been reported in several studies to be used as a fruit coating agent for tangerine oranges [17], plums [18], oranges [19], mangoes [20], and apricots [21]. The application of beeswax to enhance the quality of lime has also been reported by Kumpoun and Uthaibutra, [22] and Nasrin *et al.* [23].

Total soluble solids

The data on total soluble solids as influenced by various edible coating treatments recorded at 3rd, 6th and 9th days after storage are presented in (Table 1). The maximum TSS values (9.00, 10.20, and 11.35) at 3rd, 6th and 9th days of storage recorded in T₂ (honey bee wax 25%) and untreated fruits recorded less TSS of 3.05, 3.15, and 4.19 brix at the 3rd, 6th and 9th days, respectively. Carrillo-Lopez *et al.* [24] observed a similar thing, reporting that total soluble solids in both coated and untreated "Haden" mangoes increased during the course of storage and peaked after 16–24 days. According to Bassetto *et al.* [25], the fruit's sugar content frequently increases as it ripens because of biosynthetic processes or the breakdown of polysaccharides.

Spoilage loss

The mean data on spoilage loss (%) as influenced by various edible coating treatments recorded at the 3rd, 6th, and 9th days after storage are presented in (Table 1). The treatment (T₂) honey bee wax 25% significantly recorded the lowest spoilage loss, i.e., 0.45 and 0.69%, respectively, at the 6th and 9th days after storage. Treatment (T₁₀) control recorded maximum spoilage loss, i.e., 9.25 and 12.39% at the 6th and 9th

days after storage, respectively. Honey Bee Wax may have a delayed senescence impact by maintaining synthesis and slowing down respiration, hence increasing the firmness of fruit tissues and reducing fruit spoilage [26]. The reduced spoilage loss in the honey bee wax treatment is likely due to its ability to delay senescence by maintaining metabolic processes and reducing respiration rates. Honey bee wax also inhibits cellular disintegration [27-29].

Table 1 Effect of edible coatings on shelf life and quality of Mango (*Mangifera indica*)

Treatments	Physiological loss in weight (%)			TSS (Brix)			Spoilage loss (%)		Days to shriveling initiation		Fruit color	Shelf life (Days)
	3 rd day	6 th day	9 th day	3 rd day	6 th day	9 th day	6 th day	9 th day	9 th day	9 th day	-	
T ₁	0.21	0.35	0.93	9.07	10.56	13.35	0.75	1.18	7.50	8.53	12.56	
T ₂	0.19	0.25	0.33	9.29	10.88	14.14	0.45	0.69	10.50	9.06	16.35	
T ₃	0.27	0.46	1.00	9.00	10.20	12.35	0.90	2.50	8.50	7.22	11.25	
T ₄	4.59	8.74	15.28	9.27	10.78	14.42	8.00	9.75	6.75	6.72	10.89	
T ₅	4.17	7.46	13.35	9.02	10.33	12.44	5.75	9.00	7.25	7.15	11.20	
T ₆	4.59	7.71	14.28	9.25	10.76	13.70	4.50	8.00	7.50	8.77	12.76	
T ₇	4.11	7.23	12.89	5.15	6.25	7.39	1.44	3.73	4.57	3.17	7.01	
T ₈	4.18	7.80	13.42	3.17	4.46	4.57	3.79	5.23	5.82	4.20	8.28	
T ₉	2.11	3.83	4.75	4.29	4.84	5.80	4.25	5.39	6.63	3.78	8.05	
T ₁₀	5.13	10.72	16.42	3.05	3.15	4.19	9.25	12.39	4.27	2.07	6.56	
S.Ed.	0.09	0.32	0.37	0.15	0.15	0.33	0.98	1.05	0.82	0.09	1.20	
CD (P=0.05)	0.27	0.97	1.10	0.34	0.46	1.00	1.24	2.27	1.63	0.27	2.51	

T₁: Honey Bee Wax 10%
T₂: Honey Bee Wax 25%
T₃: Honey Bee Wax 30%
T₄: Aloe Vera Gel 10%
T₅: Aloe Vera Gel 25%

T₆: Aloe Vera Gel 30%
T₇: Gelatin 10%
T₈: Gelatin 25%
T₉: Gelatin 30%
T₁₀: Control

Days to shriveling initiation

Edible coating treatments were recorded at 3rd, 6th, and 9th after storage, as presented in (Table 1). The treatment (T₂) Honey Bee Wax 25% recorded the late initiation of shriveling, i.e., 10.50 days after storage, as compared to the rest of the treatments. An early shriveling was recorded in the fruits of treatment (T₁₀) at 4.27 days after storage. This could be because the cell wall has more contributing strength. Regardless of the amount of honey bee wax present in the tissue cell wall in vivo, a significantly higher concentration may have added strength, allowing for a delay in the cell wall's breakdown and subsequent increase in permeability. The overall effect of such an intermediary-modified cell wall structural feature may alter the morphological characteristics of the entire mango fruit [30].

Fruit colour

After the organoleptic test, the fruit color at 9th day after storage was recorded and is presented in (Table 1). The treatment (T₂) Honey Bee Wax 25% significantly recorded the most attractive fruit color and found the maximum value of fruit color, i.e., 9.06, at the 9th day after storage, and the minimum was recorded in the control (2.07). Fruit colour changes during storage are associated with the production of pigments like carotenoids. Given that the beeswax coating affected the fruit's internal gas composition and delayed the diffusion of gases, it was plausible that the delayed colour changes in the coated fruits were caused by a decrease in the activity of enzymes involved in the synthesis of carotenoids. However, the reduced colour loss in the coated fruit may have been explained by the beeswax coating's capacity to postpone the synthesis of carotenoids [31]. In a similar vein, Mladenoska [32] and Nasrin

et al. [33] noted that the appearance of strawberry and lemon fruits was enhanced by beeswax coatings containing coconut oil. The beeswax coating may act as a barrier to prevent the diffusion of oxygen and carbon dioxide, which could lead to a decrease in respiration and other metabolic processes that could alter the color of the material.

Shelf-life

The treatment (T₂) Honey Bee Wax 25% recorded a significantly maximum shelf life (16.35 days), but the treatment (T₁₀) control recorded a minimum of 6.56 days. The current findings corroborated those of Penchaiya et al. [34] applying an edible covering increased the shelf life of mango fruits.

CONCLUSION

The findings indicate that mango fruits treated with 25% honey bee wax had lower physiological weight loss, minimal TSS content, delayed shrivelling initiation, and reduced spoilage loss. The highest organoleptic scores were recorded for fruit treated with honey bee wax on the third, sixth, and ninth days of storage, which included colour, pulp colour, taste, and aroma. The promise of beeswax as an edible fruit coating substance is supported by these findings. Additionally, compared to the expensive cost of commercial coating chemicals, utilizing beeswax may have economic benefits, particularly in underdeveloped nations. Therefore, if fresh mangoes are handled carefully with the right kind of wax coat and concentration, mango supervisors can readily adapt the technology to treat fresh mangoes in order to transport them over great distances without compromising their quality.

LITERATURE CITED

1. Talcott ST, Moore JP, Lounds-Singleton AJ, Percival SS. 2005. Ripening associated phytochemical changes in mangos (*Mangifera indica*) following thermal quarantine and low-temperature storage. *Jr. Food Science* 70(5): 337-347.

2. Alam MK, Rana ZH, Islam SN. 2016. Comparison of the proximate composition, total carotenoids and total polyphenol content of nine orange-fleshed sweet potato varieties grown in Bangladesh. *Foods* 5(3):64.
3. Gomer-Lim MA. 1997. Postharvest physiology. The Mango: Botany, production and uses. Cab Intl. New York. pp 425-446.
4. Sun X, Follett P. 2024. Effect of X-ray irradiation and carnauba wax coating on quality of lime (*Citrus latifolia* Tan.) fruit. *Hort. Science* 59(5): 684690. 2024. <https://doi.org/10.21273/HORTSCI17743-24>
5. Mahfoudhi N, Chouaibi M, Hamdi S. 2014. Effectiveness of almond gum trees exudates as a novel edible coating for improving postharvest quality of tomato (*Solanum lycopersicum* L.) fruits. *Food Sci. Technol. International* 20: 33-43.
6. Kaur J, Jawandha SK, Gill PS. 2022. Effect of beeswax enriched with sodium nitroprusside coating on antioxidant properties and quality of lemon cv. PAU Baramasi Lemon-1 fruits during low temperature storage. *Jr. Fd. Proc. Preservation* 46(3): 16319.
7. Eshetu A, Ibrahim AM, Forsido SF, Kuyu CG. Effect of beeswax and chitosan treatments on quality and shelf life of selected mango (*Mangifera indica* L.) cultivars. *Heliyon* 5(1): e01116.
8. Kumpoun W, Uthaibuttra J. 2014. Shelf and storage life extension by beeswax coating for 'Pan' lime fruit from two production locations. In: III Asia Pacific Symposium on Postharvest Research, Education and Extension. pp 171-176.
9. Amin U, Khan MKI, Khan MU, Ehtasham Akram M, Pateiro M, Lorenzo JM, Maan AA. 2021. Improvement of the performance of chitosan - aloe vera coatings by adding beeswax on postharvest quality of mango fruit. *Foods* 10: 2240.
10. Ochiki S, Gesimba MR, Ngwela WJ. 2015. Effect of Aloe vera gel coating on postharvest quality and shelf life of mango (*Mangifera indica* L.) fruits Var. 'Ngowe'. *Journal of Horticulture and Forestry* 7(1): 1-7.
11. Dadzie RG, Amoah RS, Ampofo-Asiama J, Quaye B, Kizzie-Hayford N, Abano EE. 2021. Improving the storage quality of eggplant (*Solanum aethiopicum* L.) fruits using Aloe vera gel coating. *Journal of Food Technology Research* 8(2): 58-66.
12. Quirino AK, Costa JD deS, Neto AF, Costa M deS, Sánchez-Sáenz CM. 2018. Conservation of "Paluma" guavas coated with cassava starch and pectin. *DYNA* 85(204): 344-351.
13. Neeta BT, Gol V, Ramana Rao. 2014. Influences of gelatine coatings on the postharvest quality and shelf-life extension of Mango (*Mangifera indica* L.). *Fruits* 69(2): 101-115.
14. Panse VG, Sukhatme PV. 1985. *Statistical Method for Agricultural Workers*. ICAR, New Delhi, India. pp 152-161.
15. Kumpoun W, Uthaibuttra J. 2018. Shelf and storage life extension by beeswax coating for 'Pan' lime fruit from two production locations. *Acta Horticulturae* 1213: 171-176.
16. Shahid MN, Abbasi N. 2011. Effect of bee wax coatings on physiological changes in fruits of sweet orange CV Blood red. *Sarhad Journal of Agriculture* 27: 385-394.
17. Hassan Z, Lesmayati S, Qomariah R, Hasbianto A. 2014. Effects of wax coating applications and storage temperatures on the quality of tangerine citrus (*Citrus reticulata*) var. Siam Banjar. *International Food Research Journal* 21: 641-648.
18. Gunaydin S, Karaca H, Palou L, De La Fuente B, Perez-Gago MB. 2017. Effect of hydroxypropyl methylcellulose- beeswax composite edible coatings formulated with or without antifungal agents on physicochemical properties of plums during cold storage. *Journal of Food Quality* 9. 2017: Article ID 8573549
19. Babarabie M, Sardoei AS, Jamali B, Hatami M. 2024. Carnauba wax-based edible coatings retain quality enhancement of orange (*Citrus sinensis* cv. Moro) fruits during storage. *Science Reporter* 14(1): 4133.
20. Hoseini MSM, Milani JM, Motamedzadegan A, Yousefnia PH. 2023. Effect of coating using beeswax and sodium nitroprusside on chlorophyll stability and quality factors of lime during cold storage. *Fd. Sci. Tech. International* 9: 10820132231210319
21. Kaur J, Jawandha SK, Gill PS, Jan Z, Grewal SK. 2023. Composite coatings of beeswax + naphthalene acetic acid preserved fruit quality and antioxidants in stored lemon fruits. *Food Sci. Biotechnology* 33(3): 589-598.
22. Chen H, Sun Z, Yang H. 2019. Effect of carnauba wax-based coating containing glycerol monolaurate on the quality maintenance and shelf-life of Indian jujube (*Zizyphus mauritiana* L.) fruit during storage. *Sci. Horticulturae* 244: 157-164.
23. Yimenu SM, Abera S, Solomon W. 2017. Effect of bee wax and linseed oil coatings and frequency of dipping on the biochemical and organoleptic quality of fresh orange juice (*Citrus sinensis* cv. Valencia). *Journal of Postharvest Technology* 5: 17-28.
24. Carrillo-Lopez A, Ramirez-Bustamante F, Valdez-Torres JB, Rojas-Villegas R, Yahia EM. 2000. Ripening and quality changes in mango fruit as affected by coating with an edible film. *Jr. Food Quality* 23(5): 479-486.
25. Bassetto E, Jacomino AP, Pinheiro AL. 2005. Conservation of Pedro Sato guavas under treatment with 1-methylcyclopropene, Pesqui. *Agropecuária Bras* 40(5): 433-440.
26. Nunes C, Silva M, Farinha D, Sales H, Pontes R, Nunes J. 2023. Edible coatings and future trends in active food packaging- fruits' and traditional sausages' shelf life increasing. *Foods* 12(17): 3308.
27. Priya K, Thirunavookarasu N, Chidanand DV. 2023. Recent advances in edible coating of food products and its legislations: A review. *Jr. Agric. Food Research* 12: 100623.
28. Kumar JY, Singh V, Johar V. 2022. The effect of edible coating materials to enhance storage life of guava (*Psidium guajava* L.) cv. L-49. *Pharma Innovation Jr. Hort.* 11(6): 522-525.
29. Khorram F, Ramezanian A, Hosseini SMH. 2017. Effect of different edible coatings on postharvest quality of 'Kinnow' Mandarin. *Jr. Food Meas. Charact.* 11: 1827-1833.
30. Jain SK, Mukherjee S, Gupta NK. 2001. Effect of post-harvest of treatments and storage conditions on the quality of mango during storage. *Haryana Jr. Hort. Science* 30(3/4): 183-187.
31. Ullah A, Abbasi N, Shafque M, Qureshi AA. 2017. Influence of edible coatings on biochemical fruit quality and storage life of bell pepper cv. Yolo Wonder. *Journal of Food Quality* 11: 23.
32. Mladenoska I. 2012. The potential application of novel beeswax edible coatings containing coconut oil in the minimal processing of fruits. *Advanced Technologies* 1: 26-34.
33. Nasrin TAA, Rahman MA, Arfn MS, Islam MN, Ullah MA. 2020. Effect of novel coconut oil and beeswax edible coating on postharvest quality of lemon at ambient storage. *Journal of Agriculture and Food Research* 2: 19.
34. Penchaiya P, Jansasithorn R, Kanlavanarat S. 2006. Effect of 1-mcp on physiological changes in mango Nam Dokmai. IV International conference on managing quality in chains-the integrated view on fruits and vegetables quality. pp 717-722.