



Effect of *Indigofera aspalathoides* on complete Freund's adjuvant-induced arthritis in rats

B. Raj Kapoor, S. Kavimani, V. Ravichandiran, K. Sekhar, R. Senthil Kumar, M. Rupesh Kumar, M. Pradeepkumar, John Wilking Einstein & E.P. Kumar

To cite this article: B. Raj Kapoor, S. Kavimani, V. Ravichandiran, K. Sekhar, R. Senthil Kumar, M. Rupesh Kumar, M. Pradeepkumar, John Wilking Einstein & E.P. Kumar (2009) Effect of *Indigofera aspalathoides* on complete Freund's adjuvant-induced arthritis in rats, *Pharmaceutical Biology*, 47:6, 553-557, DOI: [10.1080/13880200902902489](https://doi.org/10.1080/13880200902902489)

To link to this article: <https://doi.org/10.1080/13880200902902489>



Published online: 01 Jun 2009.



Submit your article to this journal [↗](#)



Article views: 1184



View related articles [↗](#)



Citing articles: 7 View citing articles [↗](#)

RESEARCH ARTICLE

Effect of *Indigofera aspalathoides* on complete Freund's adjuvant-induced arthritis in rats

B. Rajkapoor¹, S. Kavimani², V. Ravichandiran³, K. Sekhar³, R. Senthil Kumar⁴, M. Rupesh Kumar⁵, M. Pradeepkumar¹, John Wilking Einstein¹, and E.P. Kumar¹

¹Department of Pharmacology, St. John's Pharmacy College, Bangalore, India, ²College of Pharmacy, Mother Theresa Institute of Health Sciences, Gorimedu, Pondicherry, India, ³Department of Pharmacology, Vel's College of Pharmacy, Pallavaram, Chennai, Tamil Nadu, India, ⁴Swamy Vivekanandha College of Pharmacy, Tiruchengodu, Tamil Nadu, India, and ⁵Department of Pharmacology, Bharathi College of Pharmacy, Bharathi Nagara, Mandya District, Karnataka, India

Abstract

The effect of ethanol extract of stems of *Indigofera aspalathoides* Vahl (Papilionaceae) (EIA) was evaluated for anti-arthritic activity on complete Freund's adjuvant-induced (CFA-induced) arthritis in rats. The EIA was administered orally at doses of 250 and 500 mg/kg daily for 30 days. The paw volume was measured on days 7, 14, 21 and 30. At the end of day 30, the rats were sacrificed and various biochemical parameters such as serum aspartate transaminase (AST), alanine aminotransferase (ALT), alkaline phosphatase (ALP), total cholesterol and triglycerides were estimated. Antioxidant enzymes, such as superoxide dismutase (SOD), catalase, glutathione peroxidase (GPx) and lipid peroxide (LPO) in the liver and kidney of normal, arthritic control and EIA treated rats were studied. Oral administration of EIA effectively inhibits rat paw edema in a dose-dependent manner. EIA significantly ($P < 0.01$) altered the biochemical parameters which were affected in arthritic rats. There was significant alteration in LPO, SOD, catalase, and GPx levels when compared to arthritic control rats. Our findings showed a significant anti-arthritic effect of EIA against CFA-induced arthritis in rats.

Keywords: Antioxidants; biochemical parameters; complete Freund's adjuvant-induced arthritis; *Indigofera aspalathoides*

Introduction

Rheumatoid arthritis is a chronic progressive inflammatory arthritis. Arthritis has been found to be the most prominent manifestation of rheumatoid disease. It is a generalized connective tissue disorder that involves para-articular structures such as tendons, bursae and the tendon sheath as well as extra-articular tissues such as subcutis, lymph nodes, skeletal muscle, central and peripheral nervous system and eyes.

Complete Freund's adjuvant-induced arthritis has been used as a model of sub-chronic or chronic inflammation in rats and is of considerable relevance for the study of pathophysiological and pharmacological

control of inflammatory processes, as well as the evaluation of analgesic potential or anti-inflammatory effects of drugs (Butler et al., 1992; Besson & Guilbaud, 1988). One of the reasons for the wide utilization of this model is due to the strong correlation between the efficiency of therapeutic agents in this model and in rheumatoid arthritis in humans (Besson & Guilbaud, 1988). Anti-inflammatory drugs, currently available for the treatment of joint inflammation of various kinds, have many undesirable side effects such as the induction of peptic ulcers. Therefore, plant remedies have become increasingly popular and are often preferred to synthetically derived pharmaceuticals because they have lesser side effects.

Address for Correspondence: B. Rajkapoor, Department of Pharmacology, St. John's Pharmacy College, No. 6. 9th Cross, 2nd Main Vijyanagar, Bangalore - 560 040, India. Tel.: +91-80-2330 0958; E-mail: rajkapoorb@yahoo.co.in

(Received 18 February 2008; revised 17 May 2008; accepted 17 May 2008)

ISSN 1388-0209 print/ISSN 1744-5116 online © 2009 Informa UK Ltd
DOI: 10.1080/13880200902902489

<http://www.informapharmascience.com/phb>

Indigofera aspalathoides Vahl (Papilionaceae) is a low under-shrub with wide distribution, found mostly in south India and Ceylon. It is popularly known as "sivanar vembu" in Tamil. The stem of the plant is traditionally used for various skin disorders and tumors (CSIR, 2001). In our previous studies, the stem has been found to have anti-inflammatory and anticancer activity against Ehrlich ascites carcinoma tumors and hepatocellular carcinoma (Rajkapoor et al., 2003, 2004, 2005). The phytochemical studies of the plant have revealed the presence of 5,4'-dihydroxy 6,8-dimethoxy 7-O-rhamnosyl flavones and pterocarpan (Rajkapoor et al., 2007; Selvam et al., 2004). The present study evaluated the anti-arthritic activity of *Indigofera aspalathoides* on complete Freund's adjuvant-induced arthritis in rats.

Materials and methods

Plant material and extraction

Stems of *Indigofera aspalathoides* were collected in and around Salem district in the month of December 2005 and authenticated by G. Murthy, Botanical Survey of India, Coimbatore, Tamil Nadu, India. The stems were shade-dried for one month, then pulverized. The powder was treated with petroleum ether for de-waxing and removal of chlorophyll. Later, it was packed (250 g) in a Soxhlet apparatus and subjected to hot continuous percolation (78°C) for 8 h using 450 mL of ethanol (95% v/v) as solvent. The extract was concentrated under vacuum, dried in a desiccator (yield, 4.5% w/w) and suspended in 5% gum acacia for the study.

Rats

Male Wistar rats (100–125 g) aged 6 weeks were procured from Tamil Nadu Veterinary College, Chennai, India. They were housed in standard microton boxes with standard laboratory diet (Hindustan Lever, Mumbai, India) and water *ad libitum*. The experiments were performed in accordance with the guidelines established by the European community for the care and use of laboratory rats and were approved by the Institutional Animal Ethics Committee (IAEC).

Experimental induction of arthritis

The rats were divided into four groups with each group containing six rats. The treatment schedules of rats belonging to the different groups are shown below.

Group 1: Normal (normal saline 2 mL/kg)

Group 2: Complete Freund's adjuvant (0.1 mL)

Group 3: Ethanol extract of *Indigofera aspalathoides* (250 mg/kg, per oral)

Group 4: Ethanol extract of *Indigofera aspalathoides* (500 mg/kg, per oral)

On day zero, arthritis was induced in rats belonging to groups 2, 3 and 4 by injecting 0.1 mL of complete Freund's adjuvant (Sigma, St. Louis, MO, USA) below the plantar aponeurosis of the right foot paw of the rats (Newbould, 1963). After this day, the extract was administered to the rats at the doses of 250 and 500 mg/kg, p.o. for up to 30 days. The paw volume was measured by using plethysmograph apparatus at days 7, 14, 21, and 30. At the end of the experimental period the rats were fasted overnight and then killed by cervical decapitation. Blood was collected and serum separated out. The liver and kidney were immediately removed and suspended in ice-cold saline for antioxidant studies.

Biochemical estimation

Serum was analyzed for the following biochemical parameters: serum aspartate transaminase (AST) (Reitman & Frankel, 1957), alanine aminotransferase (ALT) (Reitman & Frankel, 1957), alkaline phosphatase (Kind & King, 1954), total protein (Lowry et al., 1951), cholesterol (Wybenga & Pileggi, 1970), and triglyceride (Hawk et al., 1954). A 10% homogenate of the tissue was used for the assay of lipid peroxidation (LPO) (Devasagayam & Tarachand, 1987), superoxide dismutase (Marklund & Marklund, 1997), catalase (Sinha, 1972), and glutathione peroxidase (GPx) (Rotruck et al., 1973).

Statistical analysis

The results are expressed as mean $\pm$ SD and the statistical significance was analyzed by one way ANOVA followed by Tukey's multiple comparison test. P values < 0.05 were considered as significant.

Results and discussion

Table 1 shows the time course of edema and inhibition rate after the administration of CFA and EIA. The hind paw developed edema in the footpad. Edema value of the injected footpad significantly ($P < 0.001$) increased and reached a peak at 30 days. Administration of EIA at a doses of 250 and 500 mg/kg of body weight significantly ($P < 0.001$) inhibited the development of the swelling induced by CFA. The doses of EIA exhibited anti-inflammatory activity in a dose-dependent manner, which was maintained until the experiment was terminated on day 30 (groups 3 and 4).

As a result of arthritis induced by CFA in group 2, the level of AST, ALT and ALP were increased (Table 2). It also showed the lipid profiles following the administration of EIA at 250 and 500 mg/kg in group 3 and 4 rats, in which a significant difference in the concentration of total cholesterol and triglyceride was found.

With adjuvant treatment, an increase in lipid peroxide was seen in liver and kidney. An arthritic rat treated with EIA was found to have altered values which were closer

to that of controls. A decrease in SOD, catalase and GPx levels in liver and kidney was observed in adjuvant-induced arthritis (Tables 3 and 4). Treatment with EIA in adjuvant-induced rats altered the antioxidant levels near to normal in a dose-dependent manner.

The adjuvant produces arthritis predominantly in the joints of hind limbs, promoting significant reduction of motor activity and an increase in itching and scratching behaviors (Calvino et al., 1987). In addition, increased

Table 1. Effect of EIA on CFA-induced paw edema in rats.

Treatment	Dose (mg/kg)	Paw volume (mL)			
		Day 7	Day 14	Day 21	Day 30
Normal (normal saline)	2 mL/kg	0.21 ± 0.01	0.20 ± 0.01	0.26 ± 0.01	0.24 ± 0.01
CFA	0.1 mL	0.56 ± 0.02 ^a	0.64 ± 0.01 ^a	0.72 ± 0.02 ^a	0.74 ± 0.03 ^a
EIA	250	0.27 ± 0.01 ^{b, d}	0.32 ± 0.01 ^{a, d}	0.36 ± 0.01 ^{a, d}	0.39 ± 0.01 ^{a, d}
	500	0.24 ± 0.01 ^d	0.28 ± 0.02 ^{c, d}	0.31 ± 0.01 ^d	0.34 ± 0.01 ^{c, d}

N = 6 rats in each group.

^aP < 0.001; ^bP < 0.05; ^cP < 0.01 versus normal; ^dP < 0.001 versus CFA.

Data were analyzed by one way ANOVA followed by Tukey multiple comparison test.

Table 2. Effect of EIA on biochemical parameters against CFA-induced arthritis in rats.

Treatment	Dose (mg/kg)	AST U/L	ALT U/L	ALP U/L	Total protein mg%	Cholesterol mg%	Triglyceride mg%
Normal (normal saline)	2 mL/kg	54.5 ± 1.65	51.25 ± 1.78	177.5 ± 3.52	6.27 ± 0.34	61.75 ± 1.8	67 ± 1.36
CFA	0.1 mL	156.2 ± 3.49 ^a	67.8 ± 1.90 ^a	301 ± 5.48 ^a	7.46 ± 0.49	68.8 ± 1.15 ^b	99.2 ± 1.59 ^a
EIA	250	122.0 ± 2.64 ^{a, d}	62.0 ± 1.22 ^a	254.0 ± 7.4 ^{a, d}	6.36 ± 0.19	58.6 ± 1.27 ^{a, d}	78.4 ± 1.60 ^a
	500	98.5 ± 1.54 ^{a, d}	57.6 ± 1.08 ^{b, c}	206.0 ± 4.93 ^{c, d}	6.24 ± 0.35	55.8 ± 1.93 ^d	65.4 ± 1.23 ^d

N = 6 rats in each group.

^aP < 0.001; ^bP < 0.05; ^cP < 0.01 versus normal; ^dP < 0.001; ^eP < 0.01 versus CFA.

Data were analyzed by one way ANOVA followed by Tukey multiple comparison test.

Table 3. Effect of EIA on antioxidant status in liver of control and experimental animals.

Treatment	Dose (mg/kg)	LPO	SOD	CAT	GPx
Normal (normal saline)	2 mL/kg	141.8 ± 4.48	4.74 ± 0.03	46.56 ± 1.4	9.65 ± 1.50
CFA	0.1 mL	394.2 ± 11.94 ^a	1.91 ± 0.02 ^a	33.92 ± 2.59 ^b	5.15 ± 0.65
EIA	250	179.55 ± 3.18 ^{b, c}	2.51 ± 0.02 ^{a, c}	40.12 ± 1.12 ^d	7.12 ± 0.80 ^c
	500	158.6 ± 2.10 ^c	3.28 ± 0.06 ^{a, c}	44.16 ± 1.16 ^d	9.10 ± 0.35 ^e

N = 6 rats in each group.

^aP < 0.001; ^bP < 0.01 versus normal; ^cP < 0.001; ^dP < 0.01; ^eP < 0.05 versus CFA.

Data were analyzed by one way ANOVA followed by Tukey multiple comparison test.

LPO = μmoles of MDA/min/mg protein; GPx = μmoles of GSH oxidized/min/mg protein; SOD = Units/min/mg protein; CAT = μmole of H₂O₂ consumed/min/mg protein.

Table 4. Effect of EIA on antioxidant status in kidney of control and experimental animals.

Treatment	Dose (mg/kg)	LPO	SOD	CAT	GPx
Normal (normal saline)	2 mL/kg	83.62 ± 2.90	3.49 ± 0.46	36.12 ± 1.04	4.65 ± 0.84
CFA	0.1 mL	127.04 ± 3.94 ^a	1.88 ± 0.11 ^c	21.53 ± 1.47 ^a	2.31 ± 0.36 ^b
EIA	250	98.77 ± 1.56 ^{b, d}	2.13 ± 0.02 ^c	28.42 ± 1.08 ^b	3.42 ± 0.28
	500	87.5 ± 1.42 ^d	2.96 ± 0.14 ^f	31.64 ± 1.35 ^d	4.01 ± 0.17

N = 6 rats in each group.

^aP < 0.001; ^bP < 0.05; ^cP < 0.01 versus normal; ^dP < 0.001; ^eP < 0.01; ^fP < 0.05 versus CFA.

Data were analyzed by one way ANOVA followed by Tukey multiple comparison test.

LPO = μ moles of MDA/min/mg protein; GPx = μ moles of GSH oxidised/min/mg protein; SOD = Units/min/mg protein; CAT = μ mole of H₂O₂ consumed/min/mg protein.

sensitivity of the affected paw to pressure or flexion and extension of the inflamed joints and weight loss were observed. The arthritis observed in rats is associated with a hyperalgesia phenomenon and spontaneous behaviors, such as protection of the affected paw, evidenced by curving and/or elevation of the paw, as well as avoidance of supporting the body on the paw (Clatworthy et al., 1995). The hyperalgesia is more evident during the acute inflammatory phase, when spontaneous behaviors, indicative of painful response are more pronounced. Increased paw diameter (posterior and anterior) due to inflammation and edema was also observed (Cain et al., 1997). The initial inflammatory response was developed within hours, but more critical clinical signals emerged from day 10 and thereafter, and the alterations remain detectable for several weeks (Clopaert et al., 1982). The results of the present study indicate that the EIA exhibits anti-arthritic effects in rats with Freund's adjuvant-induced arthritis, on its acute as well as its chronic phases.

Tissue damage was assessed by measuring the activity of enzymes in the serum and in the respective organ, since liver impairment is also a feature of adjuvant arthritis (Marylatha et al., 1998). The increase in aminotransferase is due to its release from cells of the damaged organ (Rainsford, 1982). Subrata et al. (1994) reported a similar increase in aminotransferases observed in arthritic rats. Alkaline phosphatase has been reported to be present mainly in the blood vessels, pia-arachnoid and choroid plexes. Alkaline phosphatase activity has been reported to increase during the morphological and functional development of the tissues. Aminotransferases and alkaline phosphatase were significantly reduced in arthritic rats after the administration of EIA. This reducing effect may be related to their anti-inflammatory activity. Arthritis is characterized by inflammation and an increased cardiovascular risk. It was recently shown that active early rheumatoid arthritis is associated with dyslipidemia, which may partially explain the enhanced cardiovascular risk (Van Halm et al., 2007). Table 2 shows the lipid profiles following the administration of EIA at 250 and 500 mg/kg in group 3 and 4 rats, in which there was found a significant difference in the concentration of total cholesterol and triglyceride. Therefore, we can speculate that EIA may reduce the risk of inflammation-related diseases.

The increased lipid peroxide level noticed in arthritic rats in our study (group 2) may be due to its release from neutrophils and monocytes during inflammation (Greenwald & Moy, 1980). At the onset of inflammation there is a rapid fall in the total iron content of blood plasma followed by decreased deposition of iron proteins in the synovial fluid. The drop in plasma iron correlates closely with the activity of the inflammatory process. In the synovial fluid of inflamed joints, the iron

released during necrosis, might catalyze the formation of OH (hydroxyl) radicals from H_2O_2 , thus contributing increased lipid peroxidation in arthritis. From the literature reviewed, it is apparent that RA is exposed to oxidative stress and is prone to lipid peroxidation (Heliovaara et al., 1994). In the present study, the concentration of lipid peroxidation was significantly altered in arthritic rats after the administration of EIA, when compared with arthritic control rats.

Many cellular defense mechanisms are directed against the toxic effects of these radicals in the inflammatory process. SOD which converts super oxide radicals to H_2O_2 is widely distributed in cells having oxidative metabolism and is believed to protect such cells against the toxic effects of super oxide anion. Increased delivery of NADPH from the stimulated hexose monophosphate (HMP) shunt during inflammation is proposed to lead to the activation of SOD in arthritic rats. A decrease in superoxide dismutase in the liver and kidney has been observed in adjuvant-induced arthritis (Tables 3 and 4). Increased production of NADPH from HMP shunt during arthritis may cause a decrease in SOD activity (Marklund et al., 1987). This increase in enzyme activity appears to be protective against the intracellular oxygen free radicals (Kasama et al., 1988). Administration of EIA to arthritic rats caused a significant increase in SOD activity.

Catalase in the erythrocytes functions to protect Hb against oxidation. Liver has been reported to be a major site of lipid peroxide metabolism. The main function of catalase is to detoxify H_2O_2 , although catalase is significantly decreased in rheumatoid arthritis. Its concentration is very low to expect considerable protection against H_2O_2 in arthritic rats (Blake et al., 1981). Treatment of arthritic rats with EIA significantly increased the catalase enzyme levels. Glutathione peroxidase is localized in the cytoplasm and mitochondria, which catalyses the degradation of various peroxides by oxidizing glutathione with the formation of its conjugates. The increased activity of glutathione peroxidase in liver and kidney of arthritic rats, as shown by the present results, indicates the free radical defense system against oxidative stress and may help to explain the pathogenesis associated with arthritis. After EIA treatment, GPx levels were significantly decreased.

After EIA treatment, the alterations produced in arthritic rats with respect to lipid peroxidation and antioxidant concentrations were modulated to nearly normal levels. The antiperoxidative action observed in EIA treated arthritic rats might be due to the presence of compound-like flavonoids in EIA. These compounds have been shown to scavenge free radicals, including hydroxyl and superoxide anions and reduce the levels of lipid peroxidation in stress-induced rats (Jovanovic & Simic, 2000). In the present study we thus conclude that

oral administration of EIA has been shown to modulate the above biochemical changes observed in the adjuvant-induced arthritic rats. Further studies are in progress to isolate and characterize active principles.

Declaration of interest: The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

References

- Besson JM, Guilbaud G (1988): *The arthritic rat as model of clinical pain? Proceedings of the International Symposium on the Arthritic Rat as a Model of Clinical Pain?* Held in Saint-Paul De Venice, France, June 1988, International Congress Series 837. Amsterdam, Excerpta Medica, Elsevier, p. 257.
- Blake DR, Hall ND, Terby DA, Haliwell B, Gutteridge JMC (1981): Protection against superoxide and hydrogen peroxide in synovial fluid from rheumatoid patients. *Clin Sci* 61: 483-488.
- Butler SH, Godefroy F, Besson JM, Weil Fugazaa J (1992): A limited arthritic model for chronic pain studies in the rat. *Pain* 118: 73-81.
- Cain CK, Franeis JM, Plone MA, Emerich DF, Lindner MD (1997): Pain-related disability and effects of chronic morphine in the adjuvant-induced arthritis model of chronic pain. *Physiol Behav* 62: 199-205.
- Calvino B, Crepon Bernard MA, Le Bars D (1987): Parallel clinical and behavioral studies of adjuvant-induced arthritis in rat: Possible relationship with "chronic pain". *Behav Brain Res* 24: 11-29.
- Clatworthy AL, Llich PA, Castro GA, Walters ET (1995): Role of periaxonal inflammation in the development of thermal hyperalgesia and guarding behavior in a rat model of neuropathic pain. *Neurosci Lett* 184: 5-8.
- Clopaert EC, Meert T, Witle P, Schemitt P (1982): Further evidence validating adjuvant arthritis as an experimental model of chronic pain in the rat. *Life Sci* 3: 67-75.
- CSIR (2001): *Wealth of India, Raw Material, Vol V*. New Delhi, Publication and Information Directorate, Council of Scientific and Industrial Research, India, p. 176.
- Devasagayam TPA, Tarachand U (1987): Decreased lipid peroxidation in the rat kidney during gestation. *Biochem Biophys Res Commun* 56: 836-842.
- Greenwald RA, Moy WN (1980): Effect of oxygen derived free radicals on hyaluronic acid. *Arth Rheum* 23: 455-464.
- Hawk PB, Oser L, Summerson WH (1954): *Practical Physiology Chemistry*. New York, The Maples Press, 126-132.
- Heliovaara M, Knekt P, Aho K, Aaran RK, Alfthan G, Aromaa A (1994): Serum antioxidants and risk of rheumatoid arthritis. *Ann Rheum Dis* 53: 51-53.
- Jovanovic SV, Simic MG (2000): Antioxidants in nutrition. *Ann NY Acad Sci* 899: 326-334.
- Kasama T, Kobayashi K, Sekine F, Negishi M, Ide H, Takahashi T, Niwa Y (1988): Follow-up study of lipid peroxides, superoxide dismutase and glutathione peroxidase in the synovial membrane, serum and liver of young and old mice with collagen-induced arthritis. *Life Sci* 43: 1887-1896.
- Kind PR, King EJ (1954): Estimation of plasma phosphatase by determination of hydrolysed phenol with antipyrin. *J Clin Pathol* 7: 322-326.
- Lowry OH, Rosenbrough NT, Farr AL (1951): Protein measurement with Folin-phenol reagent. *J Biol Chem* 173: 265-275.
- Marklund S, Marklund G (1997): Involvement of superoxide anion radical in the auto oxidation of pyrogallol and a convenient assay for superoxide dismutase. *Eur J Biochem* 47: 469-474.
- Marklund SL, Kling PA, Nilsson S, Ohmun M (1987): Plasma extra cellular superoxide dismutase in the acute phase response induced by surgical trauma and inflammatory disorders. *Scand J Clin Lab Invest* 47: 567-570.
- Marylatha R, Geetha T, Varalakshmi P (1998): Effect of *Vernonia cinerea* flower extract in adjuvant-induced arthritis. *Gen Pharmacol* 31: 601-606.
- Newbould BB (1963): Chemotherapy of arthritis induced in rats by *Mycobacterial* adjuvant. *Br J Pharmacol* 21: 127-136.
- Rainsford, KD (1982): Adjuvant polyarthritis in rats: is this a satisfactory model for screening anti-arthritis drugs. *Agent Action* 12: 452-458.
- Rajkapoor B, Muruges N, Lalitha KG, Krishna DR (2007): Cytotoxic activity of flavone glycoside from the stem of *Indigofera aspalathoides* Vahl. *J Nat Med* 61: 80-83.
- Rajkapoor B, Anandan R, Jayakar B (2003): Anti-inflammatory activity of *Indigofera aspalathoides* in rats. *Biomedicine* 23: 118-121.
- Rajkapoor B, Jayakar B, Muruges N (2004): Antitumour activity of *Indigofera aspalathoides* on Ehrlich ascites carcinoma in mice. *Indian J Pharmacol* 36: 38-40.
- Rajkapoor B, Muruges N, Chodon D, Sakthisekaran D (2005): Chemoprevention of *N*-nitrosodiethylamine induced phenobarbitol promoted liver tumors in rat by extract of *Indigofera aspalathoides*. *Biol Pharm Bull* 28: 364-366.
- Reitman S, Frankel SA (1957): Colorimetric method for determination of serum glutamate oxaloacetate and glutamic pyruvate transaminase. *Am J Clin Pathol* 128: 56-58.
- Rotruck JT, Pope AL, Ganther HL, Swanson AB (1973): Selenium: Biochemical role as a component of glutathione peroxidase. *Science* 179: 588-590.
- Selvam C, Sanjay MJ, Gnana Oli R, Thilagavathi R, Asit K, Chakraborti, Bhutania KK (2004): A new cyclooxygenase (COX) inhibitory pterocarpan from *Indigofera aspalathoides*: Structure elucidation and determination of binding orientations in the active sites of the enzyme by molecular docking. *Tetrahedron Lett* 45: 311-4314.
- Sinha AK (1972): Colorimetric assay of catalase. *Anal Biochem* 47: 389-394.
- Subrata D, Ravishankar B, Bhausaar GC (1994): Investigation of the anti-inflammatory effects of *Paederia foetida*. *J Ethnopharmacol* 43: 31-38.
- Van Halm VP, Nielen MMJ, Nurmohamed MT, Schaardenburg DV, Stadt Van De RJ, Koning MHMT, Habibuw MR, Horst-Bruinsma IE, Dijkmansm BAC (2007): Lipid and inflammation: serial measurements of the lipid profile of blood donors who later developed rheumatoid arthritis. *Ann Rheum Dis* 66: 184-188.
- Wybenga DR, Pileggi V (1970): Direct manual determination of total cholesterol. *J Clin Chem* 16: 980-984.